

Maths skills

1. SI units

- Look up the following terms and write a few sentences about each:

Physical Quantities	Property of material / system that can be measured - has magnitude + unit
SI Units	System of 7 physical units - metre, kilogram, second, ampere, kelvin, candela, mole
Base Units	Fundamental unit - can't be defined by combinations of others
Derived Units	units derived from 7 base units e.g. $N = \frac{kg\cdot m}{s^2}$ derived base

- In physics all units can be derived from six base units. Research how the base units are defined.

Base Quantity	Base Unit	Definition (Note: you do not need to learn these definitions)
Length	metre (m)	Length of the path travelled by light in vacuum in $\frac{1}{299792458}$ s.
Mass	kilogram (kg)	By taking fixed value of h to be $6.62607015 \times 10^{-34} \text{ kg}\cdot\text{m}^2\cdot\text{s}^{-1}$ (m and s defined as in other boxes)
Time	second (s)	fixed numerical value of the caesium frequency as $9192631770 \text{ s}^{-1}$
Temperature	kelvin (K)	By taking fixed value of K to be $1.380649 \times 10^{-23} \text{ J K}^{-1}$
Current	ampere (A)	By taking fixed value of e as $1.602176634 \times 10^{-19} \text{ C}$

2. Derived units

In physics all non-base quantities are called derived quantities and are defined by equations.

E.g. (a) Define speed. (b) Define charge.

- (a) speed = distance / time (b) charge = current × time.

The units of these new quantities are derived units and are established from these same equations. So,	(b) The unit of speed = unit of distance / unit of time = m / s = ms^{-1} ('metres per second')*
(c) The unit of charge = the unit of current × the unit of time = A·s ('amp second')	*NOTE: At A level we write divided units, such as 'metres per second' as ms^{-1} not m/s.

In the SI system, many of these derived units get their own name. For example, the SI unit of charge is the coulomb (C). So we can say that one coulomb is equal to one amp second.

or $C = A \cdot s$

Any SI unit can be expressed in terms of base units. To find the base units work though the defining equations one by one, until you end up with the base units. For example, what are the base units of a Joule? This requires two steps:

- Energy (Work) = Force × distance moved, So one joule = one newton metre ($J = \text{N}\cdot\text{m}$)
- Force is defined from $F = m \cdot a$, so one newton = one kilogram metre per second squared (or $N = \text{kg}\cdot\text{m}\cdot\text{s}^{-2}$)
- Therefore, a joule = $\text{N}\cdot\text{m} = (\text{kg}\cdot\text{m}\cdot\text{s}^{-2}) \cdot \text{m} = \text{kg}\cdot\text{m}^2\cdot\text{s}^{-2}$

Complete the table below.

Try working these out rather than looking them up. You can use the earlier answers to help with the harder ones.

Derived quantity	Defining equation	Standard SI unit (if applicable)	Equivalent base units
speed	$s = d/t$	n/a	$\text{m}\cdot\text{s}^{-1}$
momentum	$p = m v$	n/a	$\text{kg}\cdot\text{m}\cdot\text{s}^{-1}$
acceleration	$a = (v - u)/t$	n/a	$\text{m}\cdot\text{s}^{-2}$
Force	$F = m a$	newton (N)	$\text{kg}\cdot\text{m}\cdot\text{s}^{-2}$
Power	$P = W/t$	watts (W)	$\text{kg}\cdot\text{m}^2\cdot\text{s}^{-3}$
frequency	frequency = 1/time period	Hertz (Hz)	s^{-1}
charge	$Q = I t$	coulomb (C)	A·s
potential difference	voltage = work/charge	Volts (V)	$\text{kg}\cdot\text{m}^2\cdot\text{s}^{-3}\cdot\text{A}^{-1}$
resistance	$R = V/I$	Ohms (Ω)	$\text{kg}\cdot\text{m}^2\cdot\text{s}^{-3}\cdot\text{A}^{-2}$

3. Prefixes

In Physics we have to deal with quantities from the very large to the very small. A prefix is something that goes in front of a unit and acts as a multiplier. This questions will give you practice at converting figures between prefixes.

Symbol	Name	What it means	How to convert
P	peta	10^{15}	1000000000000 $\downarrow \times 1000$
T	tera	10^{12}	10000000000 $\uparrow \div 1000$ $\downarrow \times 1000$
G	giga	10^9	10000000 $\uparrow \div 1000$ $\downarrow \times 1000$
M	mega	10^6	100000 $\uparrow \div 1000$ $\downarrow \times 1000$
k	kilo	10^3	1000 $\uparrow \div 1000$ $\downarrow \times 1000$
m	milli	10^{-3}	0.001 $\uparrow \div 1000$ $\downarrow \times 1000$
μ	micro	10^{-6}	0.000001 $\uparrow \div 1000$ $\downarrow \times 1000$
n	nano	10^{-9}	0.000000001 $\uparrow \div 1000$ $\downarrow \times 1000$
p	pico	10^{-12}	0.00000000001 $\uparrow \div 1000$ $\downarrow \times 1000$
f	femto	10^{-15}	0.0000000000001 $\uparrow \div 1000$ $\downarrow \times 1000$


Convert the figures into the prefixes required (give answers in standard form).

s	ms	μ s	ns	ps
134.6	1.346×10^5	1.346×10^8	1.346×10^{11}	1.346×10^{14}
96.21	9.621×10^4	9.621×10^7	9.621×10^{10}	9.621×10^{13}
0.773	7.73×10^{-2}	7.73×10^{-5}	7.73×10^{-8}	7.73×10^{-11}

5. Lines of best fit

Read the guidance on lines of best fit here: <https://www.matrix.edu.au/the-beginners-guide-to-physics-practical-skills/physics-practical-skills-part-4-how-to-draw-a-line-of-best-fit/>

Then, draw a line of best fit for each of the graphs.



4. Significant figures

Recap ideas about significant figures here: <https://www.bellevuecollege.edu/physics/resources/measurement/sigfigs/intro/a-uncert-sigfigs/>

For each value state how many significant figures it is stated to.

Value	Sig Figs	Value	Sig Figs	Value	Sig Figs
2	1	1066	4	1800.45	6
2.0	2	82.42	4	2.483×10^4	4
2.00	3	750000	2	2.483	4
0.136	3	310	2	5906.4291	8
0.34	2	3.10×10^2	3	200000	1
6717	4				

Add the values below then write the answer to the appropriate number of significant figures

7. Rearranging equations

Rearrange these equations to express them in the terms that follow:

1. $v = x/t$

a. $x = ?$

b. $t = ?$

$$x = vt \quad t = \frac{x}{v}$$

2. $F = m a$

a. $m = ?$

b. $a = ?$

$$m = \frac{F}{a} \quad a = \frac{F}{m}$$

3. $a = (v-u)/t$

a. $t = ?$

b. $v = ?$

c. $u = ?$

$$t = \frac{(v-u)}{a} \quad v = u + at \quad u = v - at$$

6. Gradients

Calculate the gradients of the graphs below = difference in y/difference in x

(think about $y=mx+c$ from maths).

4. $v^2 = u^2 + 2as$

a. $v = ?$

b. $a = ?$

c. $u = ?$

$$v = \sqrt{u^2 + 2as} \quad a = \frac{v^2 - u^2}{2s} \quad u = \sqrt{v^2 - 2as}$$

5. $s = ut + \frac{1}{2}at^2$

a. $u = ?$

b. $a = ?$

c. $t = ?$

$$u = \frac{s - \frac{1}{2}at^2}{t} \quad a = \frac{2(s - ut)}{t^2} \quad t = \sqrt{\frac{2(s - ut)}{a}}$$

6. $\frac{1}{R_{\text{tot}}} = \frac{1}{R_1} + \frac{1}{R_2}$


a. $R_{\text{tot}} = ?$

b. $R_1 = ?$


c. $R_2 = ?$

$$R_{\text{tot}} = \frac{1}{\frac{1}{R_1} + \frac{1}{R_2}}$$


$$R_1 = \frac{1}{R_{\text{tot}}} - \frac{1}{R_2}$$




$$m = \frac{128 - 103}{50 - 0} = 2.56$$



$$m = \frac{8 - 0}{5 - 0} = 2$$




$$m = \frac{300 - 50}{5 - 0} = 50$$



$$m = \frac{20 - 0}{5 - 0} = 4$$

$$m = \frac{(500 - 100)}{(336 - 65)} = 1.51$$



$$m = \frac{20 - 0}{5 - 0} = 4$$